

Chronome/chrone

**Franz Halberg,¹ Germaine Cornélissen,¹ Othild Schwartzkopff,¹
Mario Cagnoni,² Federico Perfetto² & Roberto Tarquini²**

1. University of Minnesota, Minneapolis, MN, USA
2. University of Florence, Florence, Italy

Correspondence to: Franz Halberg, University of Minnesota, Chronobiology Laboratories,
5-187 Lyon Laboratories, 420 Washington Ave. S.E., Minneapolis,
Minnesota 55455, USA. TEL: +1 612 624 6976; FAX: + 612 624 9989
E-mail: halbe001@maroon.tc.umn.edu

Submitted: January 2, 1999

Accepted: January 3, 1999

Neuroendocrinology Letters 1999; 20: 101

Copyright © Neuroendocrinology Letters 1999

“Chronome,” derived from *chronos* (time), *nomos* (rule, law) and in the case of biological chronomes, chromosome, describes features in time, just as cells characterize the spatial organization of life. The chronome complements the genome (derived from gene and chromosome). The chronome consists of 1) a partly genetic, partly developmental, partly environmentally influenced or synchronized spectrum of rhythms; 2) stochastic or deterministic chaos; 3) trends with growth, development, maturation and aging in health and/or trends with an elevation of disease risk, illness and treatment in disease; and 4) unresolved variability. The chronome is genetically coded: it is environmentally synchronized by cycles of the socio-ecologic habitat niche and it is influenced by the dynamics of the interplanetary magnetic field.

The chronome constituents, the chronones, algorithmically formulated endpoints, are inferentially statistically validated and resolved by the computer.

Chronomes and their chronones 1) quantify normalcy, allowing an individualized positive health quantification; 2) assess, by their alterations, the earliest abnormality, including the quantification of an elevated risk of developing one (or several) disease(s), *chronorisk*, by the alteration of one or several chronones; and 3) provide, by the study of underlying mechanisms, a rational basis in the search for measures aimed at the prevention of any deterioration in properly timed, mutually beneficial environmental-organismic interactions.

1. Cornélissen G., Halberg F. (1992) Broadly pertinent chronobiology methods quantify phosphate dynamics (chronome) in blood and urine. *Clin. Chem.* 38: 329–333